

Abstract Submitted
for the APR97 Meeting of
The American Physical Society

Sorting Category: B.7

A large-scale relativistic configuration-interaction calculation of the $3s^2\ ^1S_0 - 3s3p\ ^1,^3P_1$ transition energies in magnesiumlike ions¹ K. T. CHENG, M. H. CHEN, *Lawrence Livermore National Laboratory, Livermore, CA 94550* — We have calculated the $3s^2\ ^1S_0 - 3s3p\ ^1,^3P_1$ transition energies for neutral magnesium and Mg-like Ar^{6+} , Cu^{17+} , Kr^{24+} , and Mo^{30+} using the relativistic configuration-interaction (CI) method. These calculations are based on the relativistic no-pair Hamiltonian which includes Coulomb and retarded Breit interactions and employ finite B-spline basis functions. Quantum electrodynamic and mass polarization corrections are also calculated. For Mg-like ions studied here, intravalence correlations are treated exactly by saturating the basis with all configurations that arise from valence-valence excitations. Contributions from core polarization are also calculated with large-scale CI expansions by including dominant configurations from core-valence excitations. Agreement between theory and experiment are good for all the Mg-like ions studied here.

¹This work is performed under the auspices of the U.S. DOE by Lawrence Livermore National Laboratory under contract No. W-7405-ENG-48.

☒ Prefer Oral Session
☐ Prefer Poster Session

Kwok-Tsang Cheng
ktcheng@llnl.gov
Lawrence Livermore National Laboratory

Date submitted: January 24, 1997

Electronic form version 1.2